Table 1 - Final 2003 ABCs, TACs, and OFLs of Groundfish for the Western/Central/West Yakutat (W/C/WYK), Western (W), Central (W), Eastern (W) Regulatory Areas, and in the West Yakutat (W), Southeast Outside (SEO), and Gulf-Wide (W) Districts of the GOA. [Amounts are in mt]

Species	Area ¹	ABC	TAC	OFL
Pollock ² Shumagin Chirikof Kodiak WYK Subtotal	(610) (620) (630) (640) W/C/WYK	16,788 19,685 10,339 1,078 47,890	16,788 19,685 10,339 1,078 47,890	69,410
SEO Total	(650)	6,460 54,350	6,460 54,350	8,610 78,020
Pacific cod ³ Total	W C E	20,600 29,000 3,200 52,800	15,450 22,690 2,400 40,540	70,100
Flatfish ⁴ (deep- water)	W C WYK SEO	180 2,220 1,330 1,150	180 2,220 1,330 1,150	,
Total Rex sole	W C WYK	4,880 1,280 5,540 1,600	4,880 1,280 5,540 1,600	6,430
Total	SEO	1,050 9,470	1,050 9,470	12,320
Flathead sole	W C WYK SEO	16,420 20,820 2,900 1,250	2,000 5,000 2,900 1,250	
Total		41,390	11,150	51,560
Flatfish ⁵ (shallow-water)	W C WYK SEO	23,480 21,740 1,160 2,960	4,500 13,000 1,160 2,960	
Total		49,340	21,620	61,810

Table 1. (continued)

Species	Area ¹	ABC	TAC	OFL
Arrowtooth	M	17,990	8,000	
flounder	С	113,050	25 , 000	
	WYK	18,190	2,500	
	SEO	5,910	2,500	
Total		155,140	38,000	181,390
Sablefish ⁶	W	2,570	2,570	
	C	6,440	6,440	
	WYK	2,320	2,320	
	SEO	3 , 560	3 , 560	
Subtotal	E	5 , 880	5 , 880	
Total		14,890	14,890	20,020
Pacific ⁷	W	2,700	2,700	3,220
ocean	С	8,510	8,510	10,120
perch	WYK	810	810	
	SEO	1,640	1,640	
Subtotal	E			2,900
Total		13,660	13,660	16,240
Short	W	220	220	
raker/	С	840	840	
${ t rougheye}^8$	E	560	560	
Total		1,620	1,620	2,340
Other	W	90	90	
rockfish	С	550	550	
9,10	WYK	270	150	
	SEO	4,140	200	
Total		5,050	990	6,610
Northern	W	890	890	
Rockfish ^{10,12,15}	С	4,640	4,640	
	E	N/A	N/A	_
Total		5 , 530	5 , 530	6,560
Pelagic	W	510	510	
shelf	С	3,480	3,480	
${\tt rockfish}^{\tt 13}$	WYK	640	640	
	SEO	860	860	
Total		5 , 490	5,490	8,220
Thornyhead	W	360	360	
rockfish	C	840	840	
· · ·	E	800	800	
Total		2,000	2,000	3,050

Species	Area ¹	ABC	TAC	OFL
Demersal shelf rockfish ¹¹	SEO	390	390	540
Atka mackerel	GW	600	600	6 , 200
Other ^{14,15} species	GW	N/A	11,260	N/A
TOTAL ¹⁶		416,600	236,440	531,410

Regulatory areas and districts are defined at § 679.2.

- Pollock is apportioned in the Western/Central Regulatory areas among three statistical areas. During the A season, the apportionment is based upon an adjusted estimate of relative distribution of pollock biomass at 25 percent, 56 percent, and 19 percent in Statistical Areas 610, 620, and 630, respectively. During the B season, the apportionment is based on the relative distribution of pollock biomass at 25 percent, 66 percent, and 9 percent in Statistical Areas 610, 620, and 630, respectively. During the C and D seasons, the apportionment is based on the relative distribution of pollock biomass at 47 percent, 23 percent, and 30 percent in Statistical Areas 610, 620, and 630, respectively. These seasonal apportionments are shown in Table 3. In the West Yakutat and Southeast Outside Districts of the Eastern Regulatory Area, pollock is not divided into seasonal allowances.
- The annual Pacific cod TAC is apportioned 60 percent to an A season and 40 percent to a B season in the Western and Central Regulatory Areas of the GOA. Pacific cod is allocated 90 percent for processing by the inshore component and 10 percent for processing by the offshore component. Seasonal apportionments and component allocations of TAC are shown in Table 4.
- Deep water flatfish" means Dover sole, Greenland turbot, and deep sea sole.
- "Shallow water flatfish" means flatfish not including "deep water flatfish," flathead sole, rex sole, or arrowtooth flounder.
- Sablefish is allocated to trawl and hook-and-line gears (Table 2).
- "Pacific ocean perch" means <u>Sebastes</u> <u>alutus</u>.
- "Shortraker/rougheye rockfish" means $\frac{Sebastes\ borealis}{S.\ aleutianus}$ (rougheye).
- "Other rockfish" in the Western and Central Regulatory Areas and in the West Yakutat District means slope rockfish and demersal shelf rockfish. The category "other rockfish" in the Southeast Outside District means slope rockfish.
- "Slope rockfish" means <u>Sebastes aurora</u> (aurora), <u>S. melanostomus</u> (blackgill), <u>S. paucispinis</u> (bocaccio), S. <u>qoodei</u> (chilipepper), <u>S. crameri</u> (darkblotch), <u>S. elonqatus</u> (greenstriped), <u>S. varieqatus</u> (harlequin), <u>S. wilsoni</u> (pygmy), <u>S. babcocki</u> (redbanded), S. <u>proriqer</u> (redstripe), <u>S. zacentrus</u> (sharpchin), <u>S. jordani</u> (shortbelly), <u>S. brevispinis</u> (silvergrey), S. <u>diploproa</u> (splitnose), <u>S. saxicola</u> (stripetail), <u>S. miniatus</u> (vermilion), and S. <u>reedi</u> (yellowmouth). In the Eastern GOA only, "slope rockfish" also includes northern rockfish, <u>S. polyspinous</u>.
- "Demersal shelf rockfish" means <u>Sebastes pinniger</u> (canary), <u>S. nebulosus</u> (china), <u>S. caurinus</u> (copper), <u>S. maliger</u> (quillback), <u>S. helvomaculatus</u>

- (rosethorn), \underline{S} . $\underline{\text{nigrocinctus}}$ (tiger), and \underline{S} . $\underline{\text{ruberrimus}}$ (yelloweye).
- "Northern rockfish" means <u>Sebastes</u> polyspinis.
- "Pelagic shelf rockfish" means <u>Sebastes ciliatus</u> (dusky), <u>S</u>. <u>entomelas</u> (widow), and S. flavidus (yellowtail).
- (widow), and <u>S</u>. <u>flavidus</u> (yellowtail).

 "Other species" means sculpins, sharks, skates, squid, and octopus. The TAC for "other species" equals 5 percent of the TACs of assessed target species.
- N/A means not applicable.
- The total ABC is the sum of the ABCs for assessed target species.